

Tavistock and Portman E-Prints Online

JOURNAL ARTICLE

Original citation:

Kennedy, Eilis and Kumar, Ajit and Datta, Soumitra (2007) [Antipsychotic medication for childhood-onset schizophrenia](#). Schizophrenia Bulletin, 33 (5). pp. 1082-1083.
ISSN 0586-7614

© 2007 Eilis Kennedy

This version available at: <http://taviporttest.da.ulcc.ac.uk/>

Available in Tavistock and Portman E-Prints Online: Oct 2009

The Trust has developed the Repository so that users may access the clinical, academic and research work of the Trust.

Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in Tavistock and Portman E-Prints Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain. You may freely distribute the URL

[\(http://taviporttest.da.ulcc.ac.uk/\)](http://taviporttest.da.ulcc.ac.uk/) of Tavistock and Portman E-Prints Online.

This document is the published version of 'Antipsychotic medication for childhood-onset schizophrenia'. It is freely available online. You are encouraged to consult the remainder of this publication if you wish to cite from it.

Antipsychotic Medication for Childhood-Onset Schizophrenia

Eilis Kennedy^{1,2}, Ajit Kumar³, and Soumitra S. Datta⁴

²Child and Family Department, Tavistock Clinic, 120 Belsize Lane, London NW3 5BA, UK; ³Academic unit of Psychiatry, Leeds, UK; ⁴Department of Child and Adolescent Psychiatry, Royal Manchester Children's Hospital, Manchester, UK

Key words: cochrane/childhood onset/schizophrenia

Childhood-onset schizophrenia is defined as schizophrenia with onset prior to the age of 13 years. Although it is rare, those who suffer from schizophrenia at an early age appear to have a clinically severe form of the illness with a poor long-term prognosis. Antipsychotic medication is one way of managing this rare but serious mental illness. A new Cochrane review examines antipsychotic medication for childhood-onset schizophrenia.¹

Objective

To determine the effects of antipsychotic medication in the treatment of childhood-onset schizophrenia.

Search Strategy

We searched the Cochrane Schizophrenia Group Trials Register (February 2007), inspected references of all identified studies for further trials, and contacted relevant pharmaceutical companies and authors of trials for additional information.

Selection Criteria

We included all randomized clinical trials involving children and young people with a diagnosis of childhood-onset schizophrenia (ie, with a diagnosis of schizophrenia before the age of 13) comparing any antipsychotic drug with another antipsychotic or placebo.

Data Collection and Analysis

Studies were reliably selected, quality assessed, and data extracted. For homogenous dichotomous data, we calculated random effects, relative risk (RR) and its 95% con-

fidence interval (CI), and, where appropriate, number needed to treat on an intention-to-treat basis. For normal continuous data, we calculated the weighted mean difference (WMD).

Main Results

From a total of 2062 citations, 6 relevant trials were identified. These studies were small (total $N = 256$ participants) and short (all duration 12 weeks or less) and reported data for 3 comparisons (atypical vs. typical, atypical vs atypical, and typical vs. typical antipsychotic drugs), only one of which (atypical vs. typical) was to suggest differences between treatment groups.

A few results from one study favored the atypical antipsychotic clozapine over haloperidol for treating treatment-resistant childhood-onset schizophrenia ($n = 21$, WMD children's global assessment scale 17.00, CI 7.74–26.26; $n = 21$, WMD Bunney-Hamburg Psychosis Rating Scale -3.60 , CI -6.64 to -0.56). Participants on clozapine, however, were 3 times more likely to have drowsiness (1 randomized controlled trial [RCT], $n = 21$, RR 3.3, CI 1.2–8.9, number needed to harm 2, CI 2–17) and half of the children receiving clozapine in this small short study experienced neutropenia (1 RCT, $n = 21$, RR 12, CI 0.75–193; figure 1).

Reviewers' Conclusions

There are few relevant trials and, presently, little conclusive evidence regarding the effects of antipsychotic medication for those with the rare but difficult problem of very early-onset schizophrenia. Some benefits were identified in using the atypical antipsychotic clozapine compared with haloperidol, but the benefits were offset by an increased risk of serious adverse effects.

Implications for Practice

As yet there are few data to support the use of one antipsychotic medication over another in the treatment of schizophrenia with onset in childhood. The superiority of atypical antipsychotic medications over typical antipsychotic drugs is not reflected in studies using typical medications such as chlorpromazine as the comparator medication, rather than haloperidol. Routine practice may be best within the context of an evaluative trial.

¹To whom correspondence should be addressed; tel: 0207-435-7111, fax: 0207-447-3733, e-mail: EKennedy@tavi-port.nhs.uk.

Review: Antipsychotic medication for childhood-onset schizophrenia
 Comparison: 01 ATYPICAL vs TYPICAL ANTIPSYCHOTICS (only short term)
 Outcome: 14 Adverse effects: 6. Drop in the absolute neutrophil count below 1500 mm cube




Fig. 1. Clozapine vs Haloperidol—Neutropenia.

Implications for Research

Further RCTs with larger samples and longer treatment duration are needed to evaluate the effectiveness and safety of antipsychotic medication in the treatment of childhood-onset schizophrenia. There has been no significant increase in the number of trials in the last 3 decades possibly reflecting the challenges of undertaking research with this age group. Given the relative rarity of childhood-onset schizophrenia, future treatment studies will require multicenter collaboration and extensive case-finding efforts.

Acknowledgments

We would like to thank the Cochrane Schizophrenia Group for their help and support with this review. Disclosures: None.

Reference

1. Kennedy E, Kumar A, Datta SS. Antipsychotic medication for childhood-onset schizophrenia. *Cochrane Database Syst Rev*; 2007;Disc issue 3. Chichester, UK: John Wiley & sons.